

Cannabidiol protects C2C12 myotubes against cisplatin-induced atrophy by regulating oxidative stress

Running title: CBD prevents cisplatin-induced muscle atrophy in vitro

Olivier Le Bacquer¹, Phelipe Sanchez¹, Véronique Patrac¹, César Rivoirard¹, Nicolas Saroul^{1,2}, Christophe Giraudet¹, Ayhan Kocer³, Stéphane Walrand^{1,4}

1. Unité de Nutrition Humaine (UNH), Université Clermont Auvergne, INRAE, CRNH Auvergne, 63000 Clermont-Ferrand, France.
2. Department of Head and Neck Surgery, Clermont-Ferrand University Hospital, Clermont-Ferrand, France.
3. Université Clermont Auvergne, iGReD, CNRS UMR 6293, INSERM U1103, 28, place Henri Dunant, BP38, 63001 Clermont-Ferrand, France.
4. CHU Clermont-Ferrand, Service Nutrition Clinique, 63000 Clermont-Ferrand, France.

*correspondence to: Olivier Le Bacquer, Ph.D.

INRAE, UMR1019, Université Clermont Auvergne, UNH - Unité de Nutrition Humaine, CRNH Auvergne, 63000 Clermont-Ferrand, France.

Phone: +33 473178248 / fax: +33 473178222

Email: olivier.le-bacquer@inrae.fr

ORCID : 0000-0003-3805-973X

Supplemental data available at :

URL : [https://figshare.com/articles/figure/Supplemental_Figure_S1-](https://figshare.com/articles/figure/Supplemental_Figure_S1-AJPCellPhysiol_2024_pdf/25196435)

[AJPCellPhysiol_2024_pdf/25196435](https://figshare.com/articles/figure/Supplemental_Figure_S1-AJPCellPhysiol_2024_pdf/25196435)

DOI : <https://doi.org/10.6084/m9.figshare.25196435>

33 **ABSTRACT**

34 Cancer and chemotherapy induce a severe loss of muscle mass (known as cachexia), which
35 negatively impact cancer treatment and patient survival. The aim of the present study was to
36 investigate whether CBD administration may potentially antagonize the effects of cisplatin in
37 inducing muscle atrophy, using a model of myotubes in culture. Cisplatin treatment resulted
38 in a reduction of myotube diameter (15.7 ± 0.3 vs. 22.2 ± 0.5 μm , $p<0.01$) that was restored to
39 control level with $5\mu\text{M}$ CBD (20.1 ± 0.4 μM , $p<0.01$). Protein homeostasis was severely
40 altered with a $\approx 70\%$ reduction in protein synthesis ($p<0.01$) and a 2-fold increase in
41 proteolysis ($p<0.05$) in response to cisplatin. Both parameters were dose dependently restored
42 by CBD co-treatment. Cisplatin treatment was associated with increased TBARS content
43 (0.21 ± 0.03 to 0.48 ± 0.03 nmol/mg prot , $p<0.05$), catalase activity (0.24 ± 0.01 vs. 0.13 ± 0.02
44 $\text{nmol/min}/\mu\text{g prot}$, $p<0.01$), whereas CBD co-treatment normalized TBARS content to control
45 values (0.22 ± 0.01 nmol/mg prot , $p<0.01$) and reduced catalase activity (0.17 ± 0.01
46 $\text{nmol/min}/\mu\text{g prot}$, $p<0.05$). These changes were associated with increased mRNA expression
47 of GPX1, SOD1, SOD2 and CAT mRNA expression in response to cisplatin ($p<0.01$), which
48 was corrected by CBD co-treatment ($p < 0.05$). Last, cisplatin treatment increased the
49 mitochondrial protein content of NDUFB8, UQCRC2, COX4 and VDAC1 (involved in
50 mitochondrial respiration and apoptosis), and CBD co-treatment restored their expression to
51 control values. Altogether, our results demonstrated that CBD antagonizes the cisplatin-
52 induced C2C12 myotube atrophy and could be used as an adjuvant in the treatment of cancer
53 cachexia to help maintain muscle mass and improve patient quality of life.

54

55 **New and noteworthy:** In an in vitro model, cisplatin treatment led to myotube atrophy
56 associated with dysregulation of protein homeostasis and increased oxidative stress, resulting
57 in increased apoptosis. Co-treatment with cannabidiol was able to prevent this phenotype by
58 promoting protein homeostasis and reducing oxidative stress.

59

60 **KEYWORDS**

61 cachexia, endocannabinoid system, protein homeostasis, oxidative stress, mitochondrial
62 function

63

64 INTRODUCTION

65 Cachexia is a multifactorial wasting syndrome characterized by a severe involuntary loss of
66 body weight (i.e. more than 5% body weight loss in the last 12 months or less), loss of
67 skeletal muscle mass (with or without loss of fat mass), anorexia, and dysregulated energy
68 and protein metabolism (1). Prevalence of cachexia varies with cancer type, from 15% in
69 prostate cancer up to 60% in pancreatic cancer (2). Several studies have clearly demonstrated
70 that survival times are shorter in patients who have experienced weight loss than in those who
71 have not. Weight loss is not only predictive of survival but also of response to chemotherapy
72 (3). One of the mechanisms that has been advanced to explain why patients with cachexia
73 have poorer survival is their higher incidence of complications related to surgical,
74 radiotherapeutic and chemotherapeutic treatments (4). Cachexia is associated with increased
75 fatigue and frailty, reduced physical activity leading to loss of autonomy, and decreased
76 quality of life (5, 6). In some cases, the impact of cachexia is severe enough to necessitate
77 chemotherapy dose reductions, treatment postponements or permanent discontinuation, in
78 which case patients who lose weight do not get the full potential benefit of their cancer
79 therapy (4, 7). Maintaining skeletal muscle mass in cancer patients is therefore crucial to their
80 management, improved response to associated therapies, and improved quality of life. There
81 are no formal guidelines for the management of cancer-related muscle wasting, but an
82 effective strategy should aim to reduce muscle wasting in order to promote survival in
83 patients with advanced cancer (8). One straightforward way to do this is through nutrition (9).

84 Cancer treatment often involves chemotherapy, which itself contributes to the development
85 and progression of muscle atrophy and weakness in treated patients and preclinical models
86 (10-13). Chemotherapy has been found to reduce muscle mass index and strength in lung and
87 breast cancer patients (10), and cisplatin treatment in head and neck cancer patients decreased
88 muscle strength measured by chair-lift and arm flexion tests (11). This same type of muscle
89 dysfunction is also observed in preclinical models of cancer and chemotherapy. For example,
90 in mice, implantation of C26 colonic tumor or lung carcinoma led to muscle atrophy
91 associated with a decrease in mitochondrial respiration (14) and an alteration of the processes
92 regulating mitochondrial biogenesis and dynamics (fusion/fission) (15). Similarly, treatments
93 using cisplatin or doxorubicin, two widely-used chemotherapy agents, increased muscle
94 mitochondrial dysfunction by altering mitochondrial biogenesis and dynamics (16-18). These
95 data are consistent with the mitochondrial dysfunction, decreased mitochondrial content, and
96 alterations in mitochondrial dynamics that are well-documented in cancer patients (19). As a

97 result, damaged mitochondria accumulate in skeletal muscle and, in addition to being less
98 bioenergetically efficient, promote oxidative stress through increased production of reactive
99 oxygen species (ROS). In mice, administration of doxorubicin or cisplatin leads to an increase
100 in ROS production (16-18). Mitochondrial dysfunction and increased oxidative stress disrupt
101 protein turnover pathways, leading to decreased protein synthesis and increased activity of
102 muscle proteolytic systems (proteasome, autophagy) (12, 20), ultimately resulting in muscle
103 fiber atrophy (16, 17).

104 The endocannabinoid system is a major molecular system responsible for controlling
105 metabolism throughout the body, and is becoming an increasingly popular target for
106 pharmacotherapy. Endocannabinoids (EC) and phytocannabinoids are the two main
107 subclasses of cannabinoids. EC are produced by mammals, whereas phytocannabinoids,
108 including cannabidiol (CBD), are produced by plants such as *Cannabis sativa* (21). EC exert
109 their pharmacological effects via the endogenous endocannabinoid system, mainly by
110 interacting with various receptors, primarily CB1 and CB2 (21). CBD is a weak agonist of
111 CB1 and CB2, and can activate multiple cellular targets (e.g. TRPV1, PPAR γ) or inhibit (e.g.
112 GPR55) (22). The EC system is involved in numerous physiological processes, such as
113 memory, appetite, and the regulation of metabolic energy balance (21). EC exert a central
114 effect by stimulating food intake but also by modulating lipid and carbohydrate metabolism in
115 the liver, adipose tissue and skeletal muscle to favor energy accumulation (21). There is also
116 growing evidence that the EC system and CBD plays an important role in regulating
117 mitochondrial biogenesis, membrane integrity and oxidative capacity (23). Our laboratory and
118 other teams have demonstrated that the EC system also controls muscle development (24, 25)
119 and that alterations in the EC system are associated with muscle dysfunction (26-29). The use
120 of CBD could therefore hold benefit for improving the treatment of cancer and cancer-
121 induced cachexia, in particular by protecting skeletal muscle mass.

122 Recent studies have shown that CBD has antineoplastic and anti-inflammatory properties in
123 numerous in vitro models (30). Recent evidence also indicates that CBD regulates oxidative
124 activity and mitochondrial content in the myocardium by modulating the expression of several
125 markers of mitochondrial biogenesis that had been severely reduced by doxorubicin treatment
126 (13). In addition to its effects on mitochondria, CBD also has beneficial effects on skeletal
127 muscle. In mdx mice, i.e. a model of Duchenne muscular dystrophy, Iannotti *et al.* reported
128 that CBD was able to prevent loss of motor activity by promoting myotube formation and

129 reducing inflammation (27). CBD also reduces the production of ceramides (deleterious lipid
130 derivatives responsible for mitochondrial dysfunction) in high-fat diet-induced obesity (31).

131 Taken together, these data show that there is currently no treatment to prevent or reduce
132 cachexia, and that CBD could be a promising candidate compound for use as an adjuvant in
133 cancer treatment, due to its demonstrated positive effects on mitochondrial function, oxidative
134 stress, and skeletal muscle development, metabolism and. Here, we used a model of myotubes
135 in culture to investigate whether CBD treatment was able to counteract the muscle atrophy
136 induced by chemotherapy (cisplatin).

137

138 MATERIAL AND METHODS

139 **Chemicals and reagents.** Dulbecco's modified Eagle medium (DMEM) and phosphatase
140 inhibitor cocktail were purchased from Sigma-Aldrich (Saint-Quentin-Fallavier, France).
141 Fetal bovine serum, horse serum, trypsin-EDTA, PBS, and penicillin-streptomycin were
142 purchased from PAA (Pasching, Austria). Primary antibodies were obtained from the
143 following sources. Thr389-phosphorylated S6K (#34475, 1/1000) total S6K (#9202, 1/1000),
144 Ser473-phosphorylated Akt (#9271, 1/1000), total Akt (#9272, 1/1000), Ser51-
145 phosphorylated eIF2 α (#3398, 1/1000), total eIF2 α (#5324, 1/1000), Thr172-phosphorylated
146 AMPK (#2535, 1/1000), total AMPK (#2532, 1/1000), caspase 3 (#9662, 1/1000), and VDAC
147 (#4866, 1/1000) antibodies were from Cell Signaling Technology (distributed by Ozyme,
148 Saint-Quentin-en-Yvelines, France). Mouse anti-puromycin mAb (clone 12D10)
149 (#MABE343, 1/1000) was from Sigma-Aldrich (Saint-Quentin-Fallavier, France).
150 Horseradish peroxidase-conjugated secondary antibodies were from DAKO (Trappes,
151 France). GAPDH antibody (#9545, 1/5000) was from Sigma-Aldrich (Saint-Quentin-
152 Fallavier, France). Polyubiquitin antibody (#ENZ-ABS840-0100, 1/1000) was purchased
153 from Enzo Life Sciences (Villeurbanne, France). Total OXPHOS antibody (#MS604, 1/1000)
154 was purchased from Mitosciences (Eugene, Oregon, USA). CoxIV antibody (#MA5-31470,
155 1/1000) was from Thermo Fisher Scientific (Courtaboeuf, France). Cisplatin and cannabidiol
156 and the TBARS and catalase assay kits were purchased from Cayman Chemicals (distributed
157 by INTERCHIM, Montluçon, France).

158 **Cell culture and differentiation.** Mouse C2C12 myoblasts cells were purchased from the
159 ATCC (#CRL-1772, American Type Culture Collection; Manassas, VA). Myoblasts were
160 cultured in a growth medium composed of DMEM containing 4.5 g/L glucose, 2.4 g/L
161 sodium bicarbonate, 10% fetal bovine serum, 100 UI/mL penicillin, and 0.1 mg/mL
162 streptomycin, and incubated at 37°C in humidified air with 5% CO₂. The medium was
163 changed every other day to ensure growth until 90% confluence. Myotube formation was
164 induced by changing the growth medium to a differentiation medium consisting of DMEM
165 supplemented with 2% horse serum, 100 UI/mL penicillin, and 0.1 mg/mL streptomycin for 5
166 days before cell treatment. Passages between 4 and 10 were used for the experiments.

167 **Cell treatments.** For all experiments, C2C12 cells were incubated in differentiation medium.
168 Cisplatin was prepared extemporaneously in PBS to obtain a 0.5 mg/mL stock solution, then
169 administered to the cells at a final concentration of 50 μ M for 24–48 h. A stock solution of
170 CBD was prepared in ethanol and stored at -80°C. For each experiment, cells were pre-treated

171 for 2 h prior to cisplatin treatment with CBD concentrations corresponding to those used in
172 subsequent cisplatin conditions (ranging from 1 to 5 μ M of CBD). After this pre-treatment,
173 cells were incubated with cisplatin with or without a CBD concentration corresponding to the
174 pre-treatment CBD concentration. Control experiments were conducted with the equivalent
175 amount of PBS and ethanol used in the cisplatin and CBD settings. C2C12 viability was
176 estimated using the CellTiter-Glo® luminescent assay (Promega, France).

177 **C2C12 myotube morphology analysis.** Myotubes were photographed directly in the culture
178 plates without fixation, using an AxioCam ERc5s digital camera coupled to an AxioVert.A1
179 microscope and ZEN 2.3 software (Zeiss, Germany). Myotube diameter was measured from
180 three independent experiments on myotubes in each condition. Three random measurements
181 were performed along the length of each myotube (n=3 measurements/myotube) using the
182 ZEN 2.3 software, and the average of the three measures was considered as a single value.

183 **Western blotting and measurement of protein synthesis rate.** Protein synthesis was
184 assessed according to the SUNSET method. The SUNSET technique uses puromycin, which
185 incorporates into nascent polypeptide chains and terminates the elongation, resulting in an
186 accumulation of puromycin-conjugated peptides that reflects the rate of protein synthesis.
187 Briefly, C2C12 cells were incubated with 1 μ M puromycin for the last 30 min of experimental
188 treatments, then washed twice with ice-cold PBS and homogenized in ice-cold buffer (50 mM
189 HEPES pH 7.4, 150 mM NaCl, 10 mM EDTA, 10 mM NaPPi, 25 mM β -glycerophosphate,
190 100 mM NaF, 2 mM Na orthovanadate, 10% glycerol, 1% Triton X-100) containing 1%
191 protease inhibitor cocktail (Sigma-Aldrich, Saint-Quentin-Fallavier, France). Homogenates
192 were centrifuged at 13,000 \times g for 10 min at 4°C. Denatured proteins were separated by SDS-
193 PAGE and transferred to a PVDF membrane (Millipore, Molsheim, France). Immunoblots
194 were blocked with 0.1% TBS-Tween-20 containing 5% dry milk, and then probed overnight
195 at 4°C with primary antibodies. After several washes with 0.1% TBS-Tween-20, the
196 immunoblots were incubated with a horseradish peroxidase-conjugated secondary antibody
197 (DAKO, Trappes, France) for one hour at room temperature. Immune-reactive bands or whole
198 lanes were visualized by chemiluminescence (ECL Western blotting substrate, Thermo Fisher
199 Scientific, Courtaboeuf, France). escent secondary antibodies were visualized using an MF-
200 ChemiBIS 2.0 camera (Fusion Solo, Vilber Lourmat, France). Band densities were quantified
201 using MultiGauge 3.2 software (Fujifilm Corporation, FSVT, Courbevoie, France). An
202 internal control was used on each gel to normalize signal intensities between gels.

203 **RNA extraction and quantitative real-time PCR.** Total RNA was extracted using Trizol
204 reagent (Invitrogen) according to the manufacturer's instructions. RNA was quantified by
205 measuring optical density at 260 nm. The concentrations of the mRNAs corresponding to
206 genes of interest were measured by reverse transcription followed by real-time PCR using an
207 AriaMX Real-Time PCR System (Agilent, Les-Usis, France). One microgram of total RNA
208 was reverse-transcribed using SuperScript® III reverse transcriptase and a combination of
209 random hexamer and oligo-dT primers (Invitrogen). PCR amplification was performed in a 20
210 μ L total reaction volume. The real-time-PCR mixture contained 5 μ L of diluted cDNA
211 template, 10 μ L of 2x ONE Green® Fast qPCR premix (Ozyme, Saint-Cyr-l'École, France),
212 and 0.5 μ M of forward and reverse primers. The amplification profile was initiated by 3 min
213 incubation at 95°C to activate the hot-start Taq DNA Polymerase, followed by 40 cycles of
214 two steps: 95°C for 5 sec (denaturation step) and 60°C for 30 sec (annealing/extension step).
215 Relative mRNA concentrations were analyzed using the AriaMX software. Relative mRNA
216 abundance was calculated using the $2^{-\Delta\Delta CT}$ method with 18S as housekeeping gene.
217 Mitochondrial DNA (mtDNA) was quantified by measuring the ratio between the expression
218 of mitochondrial ND1 DNA and nuclear actin DNA as reference. Details of the primers used
219 in the PCR can be found in table 1.

220 **Statistical analysis.** Data are expressed as mean \pm SE. Between group differences were
221 analyzed using one-way ANOVA and Tukey's test for post hoc comparisons. Statistical
222 significance was set at $P < 0.05$ for all analyses.

223

224 **RESULTS**

225 **CBD prevents cisplatin-induced C2C12 myotube atrophy and death.** In a first set of
226 experiments, we studied whether CBD was able to prevent the atrophy of C2C12 myotubes in
227 response to cisplatin treatment. In absence of CBD, cisplatin induced myotube atrophy, as
228 exemplified by a $\approx 30\%$ reduction ($p < 0.01$) in myotube diameter (Fig. 1A,B). However, co-
229 treatment with CBD prevented myotube atrophy in a dose-dependent manner (Fig. 1A, B). As
230 muscle morphology analysis revealed a high degree of cell death in response to cisplatin (Fig.
231 1A), we measured ATP content as an index of cell viability. We observed that myotube
232 viability was marginally affected by cisplatin and CBD after 24 h of treatment (data not
233 shown) but was reduced by $\approx 30\%$ after 48 h of treatment (Fig. 1C). CBD restored myotube
234 viability in a dose-dependent manner, reaching full protection at a 3 μM concentration (Fig.
235 1C). Western blot analysis revealed that the reduction in cell viability observed in response to
236 cisplatin was associated with the appearance of the cleaved form of caspase 3 indicating
237 induction of apoptosis (Fig. 1D,E). Similar results were observed in myoblast (supplemental
238 figure). As expected, CBD treatment prevented the apoptotic process, as evidenced by the
239 dose-dependent reduction in protein expression of the cleaved-caspase 3 (Fig. 1D,E).

240

241 **CBD restores protein homeostasis in cisplatin-treated C2C12 myotubes.** To investigate
242 whether cisplatin alters protein synthesis and proteolysis and whether CBD is able to
243 counteract this effect, we incubated C2C12 myotubes for 24 h in differentiation media in the
244 presence of cisplatin and increasing concentrations of CBD, and then measured protein
245 synthesis using the SUnSET technique. As shown in Fig. 2A and B, puromycin incorporation
246 was reduced by $\approx 75\%$ ($p < 0.01$) in response to cisplatin treatment. Protein synthesis is mainly
247 controlled by the Akt/mTOR/S6K and eIF2 α signaling pathways, where anabolic conditions
248 lead to increased phosphorylation of Akt on Ser473 and S6K on Thr389 and decreased
249 phosphorylation of eIF2 α on Ser51 residue. As shown in Fig. 2A, C-E, cisplatin treatment
250 was associated with decreased phosphorylation of both Akt and S6K and increased
251 phosphorylation of eIF2 α , in agreement with the observed reduced protein synthesis. Co-
252 treatment with CBD was able to restore protein synthesis (Fig. 2A, B), which was associated
253 with a dose-dependent increase in Akt phosphorylation on Ser473 and S6K phosphorylation
254 on Thr389, and a dose-dependent decrease of Ser51 phosphorylation on eIF2 α (Fig. 2A, C-E).
255 AMPK kinase is activated by an elevated AMP/ATP ratio due to cellular and environmental
256 stress. We observed that cisplatin treatment was associated with an increase in AMPK

257 phosphorylation on Thr172 that was corrected by CBD co-treatment (Fig. 2A). To investigate
258 how cisplatin and cannabidiol impact proteolysis, we evaluated the level of protein poly-
259 ubiquitination and the mRNA expression of MAFBx and MuRF1. As shown in Fig. 3A,B,
260 cisplatin treatment increased protein polyubiquitination by $\approx 70\%$ ($p < 0.01$) and induced a ≈ 2 -
261 fold increase in mRNA expression of Atrogin1/MAFBx (Fig. 3C) and MuRF1 (Fig. 3D).
262 CBD reduced protein polyubiquitination levels in a dose-dependent manner (Fig. 3A,B), and
263 CBD treatment at a 5 μM concentration restored the mRNA expression of both
264 Atrogin1/MAFBx and MuRF1 to control levels (Fig. 3C,D).

265

266 **CBD prevents cisplatin-induced oxidative stress in C2C12 myotubes.** The toxicity of
267 cisplatin has been described as a function of DNA binding followed by single-stranded DNA
268 breaks. More recently, cisplatin has been shown to generate oxidative stress, which can also
269 contribute to its anti-tumor effects (32). To determine whether cisplatin-induced C2C12
270 myotube atrophy was associated with oxidative stress, we measured the level of TBARS, a
271 marker of lipid peroxidation, in C2C12 myotubes in response to cisplatin and CBD
272 treatments. We found a two-fold increase in TBARS content in response to 24 h cisplatin
273 treatment that was prevented by co-treatment with 5 μM CBD (Fig. 4A). Catalase is one of
274 the main enzymes responsible for the detoxification of hydrogen peroxide, a reactive oxygen
275 species. We therefore measured catalase activity in response to cisplatin and CBD treatments
276 (Fig. 4B). A 24 h cisplatin treatment induced a $\approx 90\%$ increase in catalase activity, indicating
277 severe oxidative stress (0.236 ± 0.020 vs 0.125 ± 0.013 nmol/min/ μg prot, $p < 0.01$) that was
278 partially restored by co-treatment with CBD (0.172 ± 0.010 vs 0.236 ± 0.020 nmol/min/ μg
279 prot, $p < 0.05$). We then measured the mRNA expression levels of several anti-oxidant
280 systems. As shown in Fig. 4C, mRNA expression of GPX1, SOD1, SOD2 and CAT were all
281 increased in response to 24 h cisplatin treatment. Co-treatment with 5 μM CBD was unable to
282 correct the mRNA expression levels of SOD2 and CAT (Fig. 4C) but decreased GPX1 and
283 SOD1 mRNA expression levels compared to cisplatin treatment (Fig. 4C).

284

285 **Effect of cisplatin and CBD treatment on mitochondria.** In order to analyze the early
286 effects of cisplatin and CBD on mitochondrial biogenesis and quality control, we estimated
287 mitochondrial density by measuring mtDNA (mitochondrial DNA) content and we measured
288 the mRNA expression levels of genes involved in mitochondrial biogenesis (PGC1 α), fission
289 (DRP1, FIS1), fusion (OPA1), and mitophagy (PRKN) in response to 24 h cisplatin and CBD
290 treatments. In these conditions, there were no significant changes in mtDNA content (Fig. 5A)

291 or mRNA expression levels of PGC1 α , FIS1, DRP1 and OPA1 (Fig. 5B-E), but we observed
292 a dramatic \approx 95% reduction in PRKN mRNA expression compared to controls (Fig. 5F).
293 Adding 5 μ M CBD to the cisplatin treatment did not result in any changes in mtDNA content
294 or mRNA expression of PGC1 α , OPA1, DRP1, FIS1 and PRKN compared to cisplatin
295 treatment alone (Fig. 5A-F). In a second step, we studied the effect of cisplatin and CBD
296 treatments on the expression of the different mitochondrial respiratory chain complexes. As
297 shown in Figure 6A-B, western blot quantification showed a marked increase in the content of
298 several mitochondrial proteins in C2C12 myotubes treated with 50 μ M cisplatin, with
299 significant increases in NDUFB8 (complex I, p <0.01), UQCRC2 (complex III, p <0.05) and
300 COX4 (complex IV, p <0.05) that were restored to control levels by co-treatment with CBD.
301 Finally, we studied the expression of VDAC1, which is a major mitochondrial transporter that
302 plays a key role in ATP production and is recognized as a key protein in mitochondria-
303 mediated apoptosis (33). Cisplatin treatment of C2C12 myotubes led to a 5-fold increase in
304 VDAC1 protein level, which was partly corrected by CBD treatment (Fig. 6C-D). In a final
305 step, we measured citrate synthase and COX activities. Treatment of C2C12 myotubes with
306 50 μ M cisplatin resulted in a significant \approx 15% increase in citrate synthase activity (Fig. 6E,
307 p <0.05) and a \approx 50% increase in COX activity (Fig. 6F, p =0.07) that were not corrected by
308 CBD co-treatment.

309

310 **DISCUSSION**

311 The objective of this study was to investigate whether treatment with CBD was able to
312 prevent chemotherapy-induced skeletal muscle atrophy in a cisplatin-treated model of
313 myotubes in culture. In a first set of experiments, we demonstrated that CBD was able to
314 prevent cisplatin-induced apoptosis and myotube atrophy. The increased protein synthesis and
315 decreased proteolysis induced by CBD in cisplatin-treated myotubes explain the anti-atrophic
316 effects of CBD. In a second set of experiments, we showed that cisplatin-induced atrophy was
317 associated with the induction of oxidative stress that was prevented by CBD co-treatment.
318 Finally, we observed early cisplatin-induced alterations in the expression of several
319 mitochondrial proteins involved in mitochondrial respiration and control of apoptosis, and
320 these alterations were also prevented by CBD co-treatment.

321

322 Mitochondria plays an important role in regulating many cellular functions, including ATP
323 production, generation of reactive oxygen species (ROS), and induction of apoptosis (34).
324 Cells maintain optimal mitochondrial health through several pathways, including

325 mitochondrial biogenesis, mitochondrial dynamics (fusion and fission processes shaping
326 mitochondrial morphology), and mitophagy (the process that removes defective mitochondria
327 through autophagy) (35). Alterations in mitochondrial distribution, morphology and function
328 have been reported in many conditions that lead to skeletal muscle wasting, including
329 cachexia and sarcopenia (36, 37). Here, we analyzed the mRNA expression levels of several
330 markers of mitochondrial dynamics in response to cisplatin and CBD treatments in myotubes
331 in culture. We observed no major effect of cisplatin or CBD on the mRNA expression levels
332 of genes involved in mitochondrial biogenesis (PGC1 α), fusion (Opa1), and fission (DRP1,
333 FIS1). However, there was a drastic reduction in the expression of the PRKN gene encoding
334 the Parkin protein. This could reflect inhibition of PARKIN-dependent mitophagy, leading to
335 an accumulation of dysfunctional mitochondria and, ultimately, muscle wasting (38). Previous
336 studies have shown that cisplatin (and chemotherapy in general) causes a decrease in
337 mitochondrial respiration and alterations in mitochondrial biogenesis and fusion/fission
338 processes in models of cancer or chemotherapy, but also in cancer patients (13, 16, 18, 19).
339 Our findings therefore seem to contradict the literature, but there are several possible
340 explanations for this divergence. The majority of studies characterizing the effect of cisplatin
341 on muscle atrophy were carried out *in vivo*, either in rodents or in cancer patients, and
342 treatments were carried out over periods of several days, sometimes with tumor implantation.
343 One limitation of our *in vitro* myotube model is that it does not represent the cellular
344 heterogeneity found in skeletal muscle, which consists of satellite cells, myoblasts, fibro-
345 adipogenic progenitor cells, and immune cells. Further research is needed to investigate the
346 impact of CBD on this heterogeneity and the coordinated functioning of different cell types.

347

348 In the present study, we also documented alterations in the expression of several proteins of
349 the mitochondrial electron transport chain (ETC). Cisplatin treatment was associated with
350 increased protein expression of NDUFB8 (complex I), UQCRC2 (complex III), and COX4
351 (complex IV) that were corrected by CBD, while protein expression of SDHB (complex II)
352 and ATP5A (complex V) was unaffected. In sarcopenia, deficient ETC activity is associated
353 with loss of muscle mass (39). Given that normal ETC function requires proportionally
354 balanced activities of these different complexes (40), the increased expression of NDUFB8,
355 UQCRC2 and COX4 could reflect imbalanced ETC activity and reduced energy production.
356 The reduced ATP content observed in cisplatin-treated myotubes and the increased level of
357 AMPK phosphorylation (indicating energy depletion) supports this hypothesis, but further
358 investigation is needed to fully understand the effects of cisplatin and CBD treatments on the

359 activity of the different mitochondrial complexes and on mitochondrial respiration. One of the
360 key proteins that control mitochondrial function is VDAC1, which plays a crucial role in the
361 release of ROS and in regulating apoptosis through the release of mitochondrial pro-apoptotic
362 factors such as cytochrome C and subsequent cleavage of caspase 3 (41). Several studies have
363 shown that the cytotoxic effect of cisplatin on cancer cell lines is associated with upregulation
364 of VDAC1 and excessive production of ROS (16, 42). To our knowledge, our study is the first
365 to demonstrate increased expression of VDAC1 in response to cisplatin in a model of skeletal
366 muscle in vitro, and its correction by CBD. It is still unclear how CBD counters these
367 mitochondrial alterations, but the answer may lie in its antioxidative properties.

368

369 Cisplatin is widely used to combat multiple types of cancers, but it has side effects such as
370 oxidative stress in muscle cells (12, 16). Antioxidative strategies are expected to be useful in
371 limiting oxidative stress-induced skeletal muscle damage (43). As expected, cisplatin
372 treatment resulted in increased TBARS levels (index of lipid peroxidation) together with
373 increased catalase activity and mRNA expression levels of several antioxidant systems
374 (GPX1, SOD1/2, CAT) aimed at detoxifying excessive ROS production. To our knowledge,
375 our study is the first to demonstrate the antioxidative properties of CBD in cisplatin-treated
376 myotubes in vitro. Previous studies have shown that CBD has an antioxidant function (22)
377 that comes from its ability to capture free radicals or transform them into less active forms.
378 For example, CBD was shown to directly prevent the formation of superoxide radicals in a
379 renal nephropathy model using cisplatin-treated mice (22), to reduce nitric oxide (NO) levels
380 in the liver of doxorubicin-treated mice (22), and to suppress ROS production by chelating
381 transition metal ions involved in the Fenton reaction (22). Note too that CBD modifies
382 cellular redox status by modulating both the expression and enzymatic activity of antioxidant
383 systems (22). Our results demonstrated that CBD was able to prevent the cisplatin-induced
384 accumulation of TBARS. The fact that CBD reduced the mRNA expression of antioxidant
385 systems (GPX1, SOD1/2, CAT) and catalase activity in response to cisplatin strongly
386 suggests a direct effect of CBD by capturing ROS and reducing NO production in response to
387 cisplatin, or by reducing their production by increasing the levels of antioxidants such as
388 glutathione (22).

389 On top of these direct antioxidant effects, CBD also indirectly modulates redox state by
390 interacting with several molecular targets, including the EC receptors CB1 and CB2, and
391 PPAR γ which controls the expression of antioxidant systems such as catalase and SOD2 (22).
392 CBD is not only a PPAR γ receptor agonist, but can also increase enzymatic antioxidant

393 activities (22). CBD could also improve oxidative state by modulating the activity of the CB1
394 and CB2 receptors. Indeed, CB1 activation increases ROS production whereas CB2 activation
395 decreases ROS production (44), and it has been shown that CBD is a negative allosteric
396 modulator of the CB1 receptor (45). Further studies are needed to fully understand which of
397 these targets mediate the protective effect of CBD in our *in vitro* model.

398 Oxidative stress is one of the primary causes of skeletal muscle atrophy in several patho-
399 physiological conditions, including muscle inactivity, muscular dystrophy, sarcopenia, and
400 cachexia (43). Regulation of skeletal muscle mass depends on the balance between protein
401 synthesis and degradation. Protein synthesis is mainly controlled by the Akt/mTOR
402 (mammalian target of rapamycin) and GCN2/eIF2 α pathways (46, 47). Excessive production
403 of ROS is known to activate PERK, a key endoplasmic reticulum stress transducer of the
404 unfolded protein response pathway (47) that activates eIF2 α and inhibits protein synthesis. In
405 skeletal muscle, excessive oxidative stress is also known to impair insulin signaling and Akt
406 activation upstream of mTOR, which is important for controlling both protein synthesis and
407 degradation (46). Given the drastic reduction of lipid peroxidation observed here with CBD in
408 cisplatin-treated myotubes, it is highly conceivable that CBD helps maintain protein
409 homeostasis by preventing excessive oxidative stress.

410

411 In summary, using a model of cultured C2C12 myotubes, we demonstrated that CBD
412 prevented cisplatin-induced atrophy by maintaining protein homeostasis (i.e. promoting
413 protein synthesis and limiting proteolysis) by reducing oxidative stress. In cancer patients,
414 muscle mass is predictive of survival but also of response to chemotherapy, which makes it
415 crucial to develop strategies for maintaining muscle mass in these patients. Cisplatin is an
416 antineoplastic agent that is commonly used in the treatment of solid tumors such as ovarian
417 carcinoma and in head and neck squamous cell carcinoma (HNSCC) (48). The toxicity of
418 cisplatin is due to its DNA binding followed by single-stranded DNA breaks, but also its
419 ability to generate oxidative stress in tumor cells. Consequently, some may consider that using
420 CBD for its antioxidant activity could undermine the efficacy of cisplatin treatment. However,
421 rather than reducing cisplatin toxicity, CBD was recently found to potentiate the antineoplastic
422 effect of cisplatin in a model of HNSCC (48). Taken together, the evidence suggests that CBD
423 could be used as an adjuvant in the treatment of cancer cachexia to help maintain muscle mass
424 and improve patient quality of life.

425

426 **GRANTS**

427 This study was supported by the INRAE AlimH (Alimentation Humaine) Department.

428 **CONFLICT OF INTEREST**

429 The authors declare they have no conflict of interest.

430

431 **FIGURE LEGENDS**

432 **Figure 1.** CBD prevents cisplatin-induced atrophy and apoptosis in C2C12 myotubes. **a)**
433 Representative pictures of myotube morphology at 24 h of incubation. **b)** Myotube diameter
434 of cisplatin and CBD treated cells. **c)** 48 h Myotube viability. **d)** Representative western blot
435 showing level of cleaved caspase 3 in response to 24 h treatments. **e)** Quantification of
436 cleaved caspase 3 level from (d). Results are expressed as mean \pm sem. ** $p < 0.01$ vs. CTL, \blacklozenge
437 $p < 0.05$ vs. CIS, $\blacklozenge\blacklozenge$ $p < 0.01$ vs. CIS, $\#\#$ $p < 0.01$ between CBD conditions.

438 **Figure 2.** CBD prevents cisplatin-induced decrease in protein synthesis in C2C12 myotubes.
439 **a)** Representative western blot of puromycin incorporation and the phosphorylation state of
440 Akt (Ser473-phospho-Akt), S6K (Thr389-phospho-S6K) and eIF2 α (Ser51-phospho- eIF2 α)
441 in response to 24 h treatment with cisplatin and CBD. Quantification of **(b)** puromycin
442 incorporation signal, **(c)** phospho-S6K, **(d)** phospho-Akt, and **(e)** phospho- eIF2 α levels.
443 Results are expressed as mean \pm sem. ** $p < 0.01$ vs. CTL, * $p < 0.05$ vs. CTL, \blacklozenge $p < 0.05$ vs.
444 CIS, $\blacklozenge\blacklozenge$ $p < 0.01$ vs. CIS, $\#\#$ $p < 0.01$ between CBD conditions.

445 **Figure 3.** CBD prevents cisplatin-induced proteolysis and atrogene expression. **a)**
446 Representative western blot showing the level of protein poly-ubiquitination in response to 24
447 h treatment with cisplatin and CBD. **b)** Quantification of total protein polyubiquitination from
448 (a). **c)** Real-time PCR quantification of *Atrogin/MAFbx* mRNA expression. **d)** Real-time PCR
449 quantification of *MuRF1* mRNA expression. Results are expressed as mean \pm sem. ** $p < 0.01$
450 vs. CTL, * $p < 0.05$ vs. CTL, \blacklozenge $p < 0.05$ vs. CIS.

451 **Figure 4.** CBD prevents cisplatin-induced oxidative stress. **a)** TBARS content in C2C12
452 myotubes treated for 24 h with cisplatin and CBD. **b)** Catalase activity. **c)** Real-time PCR
453 quantification of *Gpx1*, *Sod1*, *Sod2* and *Cat* mRNA expression. Results are expressed as mean
454 \pm sem. ** $p < 0.01$ vs. CTL, * $p < 0.05$ vs. CTL, \blacklozenge $p < 0.05$ vs. CIS. $\blacklozenge\blacklozenge$ $p < 0.01$ vs. CIS.

455 **Figure 5.** Effect of cisplatin and CBD co-treatment on C2C12 myotube mitochondrial
456 dynamics. **a)** Real-time PCR quantification of mtDNA expression in C2C12 myotubes treated

457 for 24 h with cisplatin and CBD. Real-time PCR quantification of *Pgc1alpha* (b), *Fis1* (c),
 458 *Drp1* (d), *Opal* (e), and *Prkn* (f) mRNA expression. Results are expressed as mean \pm sem. **
 459 $p < 0.01$ vs. CTL.

460 **Figure 6.** Effect of cisplatin and CBD on C2C12 myotube mitochondrial protein content and
 461 activity. a) Representative western blot showing the expression levels of proteins related to
 462 mitochondrial oxidative phosphorylation (OXPHOS) in C2C12 myotubes treated for 24 h
 463 with cisplatin and CBD. b) Quantification of several subunits of each mitochondrial complex
 464 from (a). c) Representative western blot showing the expression levels of VDAC1. d)
 465 Quantification of VDAC1 protein content from (c). (e) Citrate synthase activity and (f) COX
 466 activity. Results are expressed as mean \pm sem. ** $p < 0.01$ vs. CTL, * $p < 0.05$ vs. CTL, \blacklozenge
 467 $p < 0.05$ vs. CIS. $\blacklozenge\blacklozenge$ $p < 0.01$ vs. CIS.

468 **Table 1.** List of primers used for real-time qPCR

Gene name	5'- Sense primer -3'	5'- Antisense primer -3'
Actin (DNA primer)	TACAGCTTCACCACCACAGC	AAGGAAGGCTGGAAAAGAGC
Atrogin/MAFBx	AAGCTTGTGCGATGTTACCCA	CACGGATGGTCAGTGCCCTT
Cat	CCTTCAAGTTGGTTAATGCAGA	CAAGTTTTTGATGCCCTGGT
Drp1	TGCCTCAGATCGTCGTAGTG	TGACCACACCAGTTCCTCTG
Fis1	GCCTGGTTCGAAGCAAATAC	CACGGCCAGGTAGAAGACAT
Gpx1	GTGAGCCTGGGCTCCCTGCG	ACTTGAGGGGAATTCAGAATC
MuRF1	AGGTGTCAGCGCAAAGCAGT	CCTCCTTTGTCTCTTGCTG
Nd1 (DNA primer)	GGCCCCCTTCGACCTGACAGA	TAACGCGAATGGGCGGCTG
Opal	GATGACACGCTCTCCAGTGAAG	CTCGGGGCTAACAGTACAACC
Ppargc1a	GAAGTGGTGTAGCGACCAATC	AATGAGGGCAATCCGTCTTCA
Prkn	ATTCCAAACCGGATGAGTGG	TTGTCTGAGGTTGGGTGTGC
Sod1	CAGGACCTCATTTTAATCCTCAC	TGCCCAGGTCTCCAACAT
Sod2	GACCTGCCTTACGACTAT	TACTTCTCCTCGGTGACG
18S	CGGCTACCACATCCAAGGAA	GCTGGAATTACCGCGGCT

469

470

471 **REFERENCES**

472

473


- 474 1. **Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, Jatoi A, Loprinzi C,**
475 **MacDonald N, Mantovani G, Davis M, Muscaritoli M, Ottery F, Radbruch L, Ravasco P, Walsh D,**
476 **Wilcock A, Kaasa S, and Baracos VE.** Definition and classification of cancer cachexia: an international
477 consensus. *Lancet Oncol* 12: 489-495, 2011.
- 478 2. **Anker MS, Holcomb R, Muscaritoli M, von Haehling S, Haverkamp W, Jatoi A, Morley JE,**
479 **Strasser F, Landmesser U, Coats AJS, and Anker SD.** Orphan disease status of cancer cachexia in the
480 USA and in the European Union: a systematic review. *Journal of cachexia, sarcopenia and muscle* 10:
481 22-34, 2019.
- 482 3. **Pin F, Couch ME, and Bonetto A.** Preservation of muscle mass as a strategy to reduce the
483 toxic effects of cancer chemotherapy on body composition. *Curr Opin Support Palliat Care* 12: 420-
484 426, 2018.
- 485 4. **Andreyev HJ, Norman AR, Oates J, and Cunningham D.** Why do patients with weight loss
486 have a worse outcome when undergoing chemotherapy for gastrointestinal malignancies? *Eur J*
487 *Cancer* 34: 503-509, 1998.
- 488 5. **Persson C, and Glimelius B.** The relevance of weight loss for survival and quality of life in
489 patients with advanced gastrointestinal cancer treated with palliative chemotherapy. *Anticancer Res*
490 22: 3661-3668, 2002.
- 491 6. **Dahele M, Skipworth RJ, Wall L, Voss A, Preston T, and Fearon KC.** Objective physical activity
492 and self-reported quality of life in patients receiving palliative chemotherapy. *J Pain Symptom*
493 *Manage* 33: 676-685, 2007.
- 494 7. **Ross PJ, Ashley S, Norton A, Priest K, Waters JS, Eisen T, Smith IE, and O'Brien ME.** Do
495 patients with weight loss have a worse outcome when undergoing chemotherapy for lung cancers?
496 *British journal of cancer* 90: 1905-1911, 2004.
- 497 8. **Siddiqui JA, Pothuraju R, Jain M, Batra SK, and Nasser MW.** Advances in cancer cachexia:
498 Intersection between affected organs, mediators, and pharmacological interventions. *Biochim*
499 *Biophys Acta Rev Cancer* 1873: 188359, 2020.
- 500 9. **Roeland EJ, Bohlke K, Baracos VE, Bruera E, Del Fabbro E, Dixon S, Fallon M, Herrstedt J, Lau**
501 **H, Platek M, Rugo HS, Schnipper HH, Smith TJ, Tan W, and Loprinzi CL.** Management of Cancer
502 Cachexia: ASCO Guideline. *J Clin Oncol* 38: 2438-2453, 2020.
- 503 10. **Klassen O, Schmidt ME, Ulrich CM, Schneeweiss A, Potthoff K, Steindorf K, and Wiskemann**
504 **J.** Muscle strength in breast cancer patients receiving different treatment regimes. *Journal of*
505 *cachexia, sarcopenia and muscle* 8: 305-316, 2017.
- 506 11. **Lin KY, Cheng HC, Yen CJ, Hung CH, Huang YT, Yang HL, Cheng WT, and Tsai KL.** Effects of
507 Exercise in Patients Undergoing Chemotherapy for Head and Neck Cancer: A Pilot Randomized
508 Controlled Trial. *Int J Environ Res Public Health* 18: 2021.
- 509 12. **Sakai H, Sagara A, Arakawa K, Sugiyama R, Hirosaki A, Takase K, Jo A, Sato K, Chiba Y,**
510 **Yamazaki M, Matoba M, and Narita M.** Mechanisms of cisplatin-induced muscle atrophy. *Toxicol*
511 *Appl Pharmacol* 278: 190-199, 2014.
- 512 13. **Hao E, Mukhopadhyay P, Cao Z, Erdelyi K, Holovac E, Liaudet L, Lee WS, Hasko G,**
513 **Mechoulam R, and Pacher P.** Cannabidiol Protects against Doxorubicin-Induced Cardiomyopathy by
514 Modulating Mitochondrial Function and Biogenesis. *Mol Med* 21: 38-45, 2015.
- 515 14. **Halle JL, Pena GS, Paez HG, Castro AJ, Rossiter HB, Visavadiya NP, Whitehurst MA, and**
516 **Khamoui AV.** Tissue-specific dysregulation of mitochondrial respiratory capacity and coupling control
517 in colon-26 tumor-induced cachexia. *American journal of physiology Regulatory, integrative and*
518 *comparative physiology* 317: R68-R82, 2019.

- 519 15. **Brown JL, Rosa-Caldwell ME, Lee DE, Blackwell TA, Brown LA, Perry RA, Haynie WS, Hardee**
520 **JP, Carson JA, Wiggs MP, Washington TA, and Greene NP.** Mitochondrial degeneration precedes the
521 development of muscle atrophy in progression of cancer cachexia in tumour-bearing mice. *Journal of*
522 *cachexia, sarcopenia and muscle* 8: 926-938, 2017.
- 523 16. **Conte E, Bresciani E, Rizzi L, Cappellari O, De Luca A, Torsello A, and Liantonio A.** Cisplatin-
524 Induced Skeletal Muscle Dysfunction: Mechanisms and Counteracting Therapeutic Strategies.
525 *International journal of molecular sciences* 21: 2020.
- 526 17. **Gilliam LA, Moylan JS, Patterson EW, Smith JD, Wilson AS, Rabbani Z, and Reid MB.**
527 Doxorubicin acts via mitochondrial ROS to stimulate catabolism in C2C12 myotubes. *American*
528 *journal of physiology Cell physiology* 302: C195-202, 2012.
- 529 18. **Min K, Kwon OS, Smuder AJ, Wiggs MP, Sollanek KJ, Christou DD, Yoo JK, Hwang MH, Szeto**
530 **HH, Kavazis AN, and Powers SK.** Increased mitochondrial emission of reactive oxygen species and
531 calpain activation are required for doxorubicin-induced cardiac and skeletal muscle myopathy. *The*
532 *Journal of physiology* 593: 2017-2036, 2015.
- 533 19. **Argiles JM, Lopez-Soriano FJ, and Busquets S.** Muscle wasting in cancer: the role of
534 mitochondria. *Current opinion in clinical nutrition and metabolic care* 18: 221-225, 2015.
- 535 20. **Bowen TS, Schuler G, and Adams V.** Skeletal muscle wasting in cachexia and sarcopenia:
536 molecular pathophysiology and impact of exercise training. *Journal of cachexia, sarcopenia and*
537 *muscle* 6: 197-207, 2015.
- 538 21. **Mazier W, Saucisse N, Gatta-Cherifi B, and Cota D.** The Endocannabinoid System: Pivotal
539 Orchestrator of Obesity and Metabolic Disease. *Trends in endocrinology and metabolism: TEM* 26:
540 524-537, 2015.
- 541 22. **Atalay S, Jarocka-Karpowicz I, and Skrzydlewska E.** Antioxidative and Anti-Inflammatory
542 Properties of Cannabidiol. *Antioxidants (Basel)* 9: 2019.
- 543 23. **Lipina C, Irving AJ, and Hundal HS.** Mitochondria: a possible nexus for the regulation of
544 energy homeostasis by the endocannabinoid system? *American journal of physiology Endocrinology*
545 *and metabolism* 307: E1-13, 2014.
- 546 24. **Iannotti FA, Silvestri C, Mazzarella E, Martella A, Calvigioni D, Piscitelli F, Ambrosino P,**
547 **Petrosino S, Czifra G, Biro T, Harkany T, Tagliatela M, and Di Marzo V.** The endocannabinoid 2-AG
548 controls skeletal muscle cell differentiation via CB1 receptor-dependent inhibition of Kv7 channels.
549 *Proceedings of the National Academy of Sciences of the United States of America* 111: E2472-2481,
550 2014.
- 551 25. **Le Bacquer O, Lanchais K, Combe K, Van Den Berghe L, and Walrand S.** Acute rimonabant
552 treatment promotes protein synthesis in C2C12 myotubes through a CB1-independent mechanism.
553 *Journal of cellular physiology* 236: 2669-2683, 2021.
- 554 26. **Iannotti FA, Pagano E, Guardiola O, Adinolfi S, Saccone V, Consalvi S, Piscitelli F, Gazerro E,**
555 **Busetto G, Carrella D, Capasso R, Puri PL, Minchiotti G, and Di Marzo V.** Genetic and
556 pharmacological regulation of the endocannabinoid CB1 receptor in Duchenne muscular dystrophy.
557 *Nature communications* 9: 3950, 2018.
- 558 27. **Iannotti FA, Pagano E, Moriello AS, Alvino FG, Sorrentino NC, D'Orsi L, Gazerro E, Capasso**
559 **R, De Leonibus E, De Petrocellis L, and Di Marzo V.** Effects of non-euphoric plant cannabinoids on
560 muscle quality and performance of dystrophic mdx mice. *British journal of pharmacology* 2018.
- 561 28. **Le Bacquer O, Salles J, Piscitelli F, Sanchez P, Martin V, Montaurier C, Di Marzo V, and**
562 **Walrand S.** Alterations of the endocannabinoid system and circulating and peripheral tissue levels of
563 endocannabinoids in sarcopenic rats. *Journal of cachexia, sarcopenia and muscle* 13: 662-676, 2022.
- 564 29. **Fajardo L, Sanchez P, Salles J, Rigaudiere JP, Patrac V, Caspar-Bauguil S, Bergoglio C, Moro**
565 **C, Walrand S, and Le Bacquer O.** Inhibition of the endocannabinoid system reverses obese
566 phenotype in aged mice and partly restores skeletal muscle function. *American journal of physiology*
567 *Endocrinology and metabolism* 2023.
- 568 30. **Kis B, Ifrim FC, Buda V, Avram S, Pavel IZ, Antal D, Paunescu V, Dehelean CA, Ardelean F,**
569 **Diaconeasa Z, Soica C, and Danciu C.** Cannabidiol-from Plant to Human Body: A Promising Bioactive
570 Molecule with Multi-Target Effects in Cancer. *International journal of molecular sciences* 20: 2019.


- 571 31. **Bielawiec P, Harasim-Symbor E, Konstanynowicz-Nowicka K, Sztolsztener K, and**
572 **Chabowski A.** Chronic Cannabidiol Administration Attenuates Skeletal Muscle De Novo Ceramide
573 Synthesis Pathway and Related Metabolic Effects in a Rat Model of High-Fat Diet-Induced Obesity.
574 *Biomolecules* 10: 2020.
- 575 32. **Yu W, Chen Y, Dubrulle J, Stossi F, Putluri V, Sreekumar A, Putluri N, Baluya D, Lai SY, and**
576 **Sandulache VC.** Cisplatin generates oxidative stress which is accompanied by rapid shifts in central
577 carbon metabolism. *Scientific reports* 8: 4306, 2018.
- 578 33. **Shoshan-Barmatz V, Shteinifer-Kuzmine A, and Verma A.** VDAC1 at the Intersection of Cell
579 Metabolism, Apoptosis, and Diseases. *Biomolecules* 10: 2020.
- 580 34. **Pfanner N, Warscheid B, and Wiedemann N.** Mitochondrial proteins: from biogenesis to
581 functional networks. *Nature reviews Molecular cell biology* 20: 267-284, 2019.
- 582 35. **Leduc-Gaudet JP, Hussain SNA, Barreiro E, and Gouspillou G.** Mitochondrial Dynamics and
583 Mitophagy in Skeletal Muscle Health and Aging. *International journal of molecular sciences* 22: 2021.
- 584 36. **Carson JA, Hardee JP, and VanderVeen BN.** The emerging role of skeletal muscle oxidative
585 metabolism as a biological target and cellular regulator of cancer-induced muscle wasting. *Seminars*
586 *in cell & developmental biology* 54: 53-67, 2016.
- 587 37. **Chabi B, Ljubicic V, Menzies KJ, Huang JH, Saleem A, and Hood DA.** Mitochondrial function
588 and apoptotic susceptibility in aging skeletal muscle. *Aging cell* 7: 2-12, 2008.
- 589 38. **Peker N, Sharma M, and Kambadur R.** Parkin deficiency exacerbates fasting-induced skeletal
590 muscle wasting in mice. *NPJ Parkinsons Dis* 8: 159, 2022.
- 591 39. **Bua EA, McKiernan SH, Wanagat J, McKenzie D, and Aiken JM.** Mitochondrial abnormalities
592 are more frequent in muscles undergoing sarcopenia. *J Appl Physiol (1985)* 92: 2617-2624, 2002.
- 593 40. **Miro O, Casademont J, Casals E, Perea M, Urbano-Marquez A, Rustin P, and Cardellach F.**
594 Aging is associated with increased lipid peroxidation in human hearts, but not with mitochondrial
595 respiratory chain enzyme defects. *Cardiovasc Res* 47: 624-631, 2000.
- 596 41. **Camara AKS, Zhou Y, Wen PC, Tajkhorshid E, and Kwok WM.** Mitochondrial VDAC1: A Key
597 Gatekeeper as Potential Therapeutic Target. *Frontiers in physiology* 8: 460, 2017.
- 598 42. **Luo L, Xiong Y, Jiang N, Zhu X, Wang Y, Lv Y, and Xie Y.** VDAC1 as a target in cisplatin anti-
599 tumor activity through promoting mitochondria fusion. *Biochemical and biophysical research*
600 *communications* 560: 52-58, 2021.
- 601 43. **Lian D, Chen MM, Wu H, Deng S, and Hu X.** The Role of Oxidative Stress in Skeletal Muscle
602 Myogenesis and Muscle Disease. *Antioxidants (Basel)* 11: 2022.
- 603 44. **Han KH, Lim S, Ryu J, Lee CW, Kim Y, Kang JH, Kang SS, Ahn YK, Park CS, and Kim JJ.** CB1 and
604 CB2 cannabinoid receptors differentially regulate the production of reactive oxygen species by
605 macrophages. *Cardiovasc Res* 84: 378-386, 2009.
- 606 45. **Laprairie RB, Bagher AM, Kelly ME, and Denovan-Wright EM.** Cannabidiol is a negative
607 allosteric modulator of the cannabinoid CB1 receptor. *British journal of pharmacology* 172: 4790-
608 4805, 2015.
- 609 46. **Saxton RA, and Sabatini DM.** mTOR Signaling in Growth, Metabolism, and Disease. *Cell* 168:
610 960-976, 2017.
- 611 47. **Donnelly N, Gorman AM, Gupta S, and Samali A.** The eIF2alpha kinases: their structures and
612 functions. *Cellular and molecular life sciences : CMLS* 70: 3493-3511, 2013.
- 613 48. **Go YY, Kim SR, Kim DY, Chae SW, and Song JJ.** Cannabidiol enhances cytotoxicity of anti-
614 cancer drugs in human head and neck squamous cell carcinoma. *Scientific reports* 10: 20622, 2020.
- 615

Figure 1


A.




B.



C.



D.



E.






Figure 2


A.




B.



C.



D.



E.






Figure 3


A.



B.



C.



D.






Figure 4

A.



B.



C.




Figure 5






Figure 6



Effect of CBD on cisplatin-induced skeletal muscle atrophy in vitro



CBD prevents cisplatin-induced muscle atrophy in vitro